

Automorphic Representations of $SL(2, \mathbb{R})$ and Quantization of Fields

Do Ngoc Diep¹, Do Thi Phuong Quynh²

¹Institute of Mathematics, Vietnam Academy of Science and Technology, 18 Hoang, Quoc Viet road, Cau Giay district, 10307 Hanoi, Vietnam

²Medicine University College, Thai Nguyen University, Thai Nguyen City, Vietnam

Abstract: In this paper we make a clear relationship between the automorphic representations and the quantization through the Geometric Langlands Correspondence. We observe that the discrete series representation are realized in the sum of eigenspaces of Cartan generator, and then present the automorphic representations in form of induced representations with inducing quantum bundle over a Riemann surface and then use the loop group representation construction to realize the automorphic representations. The Langlands picture of automorphic representations is precised by using the Poisson summation formula.

Keywords: Trace formula; orbital integral; transfer; endoscopy, quantization

I. INTRODUCTION

The representation theory of $SL(2, \mathbb{R})$ is well-known in the Bargman classification: every irreducible unitary/nonunitary representation is unitarily/nonunitarily equivalent to one of representations in the five series:

- the principal continuous series representations (π_s, \mathcal{P}_s)
- the discrete series representations $(\pi_k^\pm, \mathcal{D}_n), n \in \mathbb{N}, n \neq 0$
- the limits of discrete series representations $(\pi_0^\pm, \mathcal{D}_\pm)$,
- the complementary series representations $(\pi_s, \mathcal{C}_s), 0 < s < 1$,
- the trivial one dimensional representation 1,
- the nonunitary finite dimensional representations V_k .

Looking at a Fuchsian discrete subgroup Γ of type I, i.e.

$$\Gamma \subseteq SL(2, \mathbb{Z}), \text{vol}(\Gamma \backslash SL(2, \mathbb{R})) < +\infty$$

One decomposes the cuspidal parabolic part $L_{cusp}^2(\Gamma \backslash SL(2, \mathbb{R}))$ of $L^2(\Gamma \backslash SL(2, \mathbb{R}))$, that is consisting of the so called *automorphic representations* in subspaces of automorphic forms, of which each irreducible component with *multiplicity* equal to the dimension of the space of modular forms on the upper Poincaré half-plane.

$$\mathbb{H} = \{z \in \mathbb{C} \mid \Im(z) > 0\}.$$

There are a lot of studies concerning the automorphic forms and automorphic representations. Most of them realize the representations as some induced ones. Therefore some *clear geometric realization* of these representations should present some interests.

In this paper, we use the *ideas of geometric quantization* to realize the automorphic representations in form of some Fock representations of loop algebras, see Theorem 3.7, below. In order to do this, we first present the action of the group $SL(2, \mathbb{R})$ in the induced representation of discrete series as the action of some loop algebra/group in the highest/lowest weight representations.

¹ Corresponding Author: dndiep@math.ac.vn

The representations are uniquely up to equivalence defined by the character, which are defined as some distribution function. Beside of the main goal we make also some clear presentation of automorphic representations in this context, see Theorems 3.2, 3.3. In the theorem 6.4 we expose the trace formula using both the spectral side and geometric side.

The authors thank the referee for his careful reading, the helpful remarks and comments, improving the paper.

II. ENDOSCOPY GROUPS FOR $SL(2, \mathbb{R})$

We introduce in this section the basic notions and notations concerning $SL(2, \mathbb{R})$ many of which are folklore or well-known but we collect all together in order to fix a consistent system of our notations.

The unimodular group $G = SL(2, \mathbb{R})$ is the matrix group

$$SL(2, \mathbb{R}) = \left\{ g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \middle| a, b, c, d \in \mathbb{R}, \det g = 1 \right\}$$

The group has finite center $\mathbb{Z}/2\mathbb{Z}$. Its complexified group $G_{\mathbb{C}} = SL(2, \mathbb{C})$. The unique maximal compact subgroup K of G is

$$K = \left\{ k(\theta) = \pm \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \middle| \theta \in [0, 2\pi) \right\}$$

The group is simple with the only nontrivial parabolic subgroup, which is minimal and therefore is a unique split Borel subgroup

$$B = \left\{ b = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \middle| a, b, d \in \mathbb{R}, ad = 1 \right\}$$

The Borel group B is decomposed into semidirect product of its unipotent radical N and a maximal split torus $T \cong \mathbb{R}_{+}^{*}$ and a compact subgroup $M = \{\pm 1\}$. It is well-known the Cartan decomposition $G = B \rtimes K = BK$

$$\begin{aligned} \begin{pmatrix} a & b \\ c & d \end{pmatrix} &= \begin{pmatrix} y^{1/2} & y^{-1/2}x \\ 0 & y^{-1/2} \end{pmatrix} \begin{pmatrix} \pm \cos \theta & \pm \sin \theta \\ \mp \sin \theta & \pm \cos \theta \end{pmatrix} \\ &= \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} y^{1/2} & 0 \\ 0 & y^{-1/2} \end{pmatrix} \begin{pmatrix} \pm \cos \theta & \pm \sin \theta \\ \mp \sin \theta & \pm \cos \theta \end{pmatrix} \end{aligned}$$

and it is easy to compute

$$\begin{aligned} y &= \frac{1}{c^2 + d^2} \\ \cos \theta &= \pm y^{1/2}d, \text{ or } \theta = \arccos \frac{d}{\sqrt{c^2 + d^2}} \\ \pm y^{1/2} \sin \theta &= \pm y^{-1/2}xc \cos \theta = b \text{ or } x = \pm \frac{(b+cy)}{d}, \end{aligned}$$

and the Langlands decomposition of the Borel subgroup $B = MAN$. The Lie algebra of $G = SL(2, \mathbb{R})$ is $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{R}) = \langle H, X, Y \rangle_{\mathbb{R}}$

where

$$H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, Y = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix},$$

satisfying the Cartan commutation relations

$$[H, X] = 2X, [H, Y] = -2Y, [X, Y] = H.$$

The Lie algebra of A is $\alpha = \langle H \rangle_{\mathbb{R}}$, the Lie algebra of N is $n = \langle X \rangle_{\mathbb{R}}$ The Lie algebra of B is

$$b = \alpha \oplus n = \langle H, X \rangle_{\mathbb{R}}$$

The complex Cartan subalgebra of \mathfrak{g} is a complex subalgebra

$$h = \langle H \rangle_{\mathbb{C}} \subset \mathfrak{g}_{\mathbb{C}},$$

which is coincided with its normalizer in $\mathfrak{g}_{\mathbb{C}}$. The corresponding subgroup of G such that its Lie algebra is a Cartan subalgebra, is called a Cartan subgroup. The root system of (\mathfrak{g}, h) is of rank 1 and is

$$\Sigma = \{\pm\alpha\}, \alpha = (1, -1) \in \mathbb{Z}(1, -1) \subset \mathbb{R}(1, -1).$$

There is only one positive root $\alpha = (1, -1)$, which is simple. There is a compact Cartan group $T = K = O(2)$ The coroot vector is $H_{\alpha} = (1, -1)$ and the split Cartan subgroup of B is $H = \mathbb{Z}/2\mathbb{Z} \times \mathbb{R}_{+}^* \cong \mathbb{R}^*$

Definition 2.1. An endoscopy subgroup of $G = SL(2, \mathbb{R})$ is the connected component of identity in the centralizer of a regular semisimple element of G .

Proposition 2.2. The only possible endoscopy subgroups of $G = SL(2, \mathbb{R})$, are itself or $SO(2)$.
Proof.

The regular semisimple elements of $SL(2, \mathbb{R})$, are of the form $g = \text{diag}(\lambda_1, \lambda_2)$, $\lambda_1\lambda_2 = 1$. If $\lambda_1 \neq \lambda_2$, the centralizer of g is the center $C(G) = \{\pm 1\}$ of the group $SL(2, \mathbb{R})$. If $\lambda_1 = \lambda_2$ and they are real, the centralizer of g is the group $SL(2, \mathbb{R})$, itself. If they are complex and their arguments are opposite, we have $g = \pm \text{diag}(e^{i\theta}, e^{-i\theta})$. In this case the connected component of identity of the centralizer is $SO(2)$. The endoscopy groups $SL(2, \mathbb{R})$, itself or the center $\{\pm 1\}$ are trivial and the only nontrivial endoscopy group is $SO(2)$.

III. AUTOMORPHIC REPRESENTATIONS

In this section we make clear the construction of automorphic representations. The following lemma is well-known.

Lemma 3.1. There is a one-to-one correspondence between any irreducible $2n - 1$ dimensional representations of $SO(3)$ and the n dimensional representations of $SO(2)$.

Proof: There is a short exact sequence

$$1 \longrightarrow \{\pm 1\} \longrightarrow SU(2) \longrightarrow SO(3) \longrightarrow 1 \quad (3.1)$$

The characters of the $2n - 1$, $n = 1, 2, \dots$ dimensional representation of $SO(3)$ is

$$\chi_{2n-1}(k(\theta)) = \frac{\sin(2n-1)\theta}{\sin\theta} \quad (3.2)$$

where

$$k(\theta) = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

It is well-known the Iwasawa decomposition ANK : every element $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ has a unique decomposition of form

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} y^{1/2} & 0 \\ 0 & y^{-1/2} \end{pmatrix} \begin{pmatrix} 1 & y^{-1/2}x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \quad (3.3)$$

of $SL_2(\mathbb{R})$, where

$$N = \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \right\} \quad (3.4)$$

is the unipotent radical of the Borel subgroup,

$$A = \left\{ \begin{pmatrix} y^{1/2} & 0 \\ 0 & y^{-1/2} \end{pmatrix} \right\} \quad (3.5)$$

is the split torus in the Cartan subgroup, and

$$K = \left\{ \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \right\} \quad (3.6)$$

is the maximal compact subgroup.

To each modular form $f \in Sk(\Gamma)$ of weight k on the Poincaré plane $\mathbb{H} = \text{SL}(2, \mathbb{R})/\text{SO}(2)$, we associate the automorphic form $\phi f \in Ak(\text{SL}(2, \mathbb{R}))$

$$\varphi_f \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) = y^{k/2} e^{ik\theta} f(x + iy), \quad (3.7)$$

Where x, y, θ are from the Iwasawa decomposition (3.3).

The discrete series representations are realized on the function on $L^2(\mathbb{H}, \mu_k)$, $\mu_k = y^k \frac{dx dy}{y^2}$ of weight k modular form by the formula

$$\pi_k \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) f(z) = (cz + d)^{-k} f(z). \quad (3.8)$$

Denote by D_k the discrete series representation of weight k . The cuspidal automorphic representations are realized in the space $L^2_{\text{cusp}}(\Gamma \backslash \text{SL}(2, \mathbb{R}))$ of automorphic forms

Theorem 3.2. *The set of intertwining $\text{SL}(2, \mathbb{R})$ homomorphisms from the set of discrete series representations to the set $L^2_{\text{cusp}}(\Gamma \backslash \text{SL}(2, \mathbb{R}))$ of the automorphic representations of $\text{SL}(2, \mathbb{R})$ is equal to the set $Sk(\Gamma)$ of modular forms, i.e.*

$$\text{Hom}_{\text{SL}(2, \mathbb{R})}(D_k, L^2_{\text{cusp}}(\Gamma \backslash \text{SL}(2, \mathbb{R}))) = S_k(\Gamma) \quad (3.9)$$

Proof: The theorem is well-known in literature, see for example A. Borel [B]. Starting from some intertwining operator $A \in \text{Hom}_{\text{SL}(2, \mathbb{R})}(D_k, L^2_{\text{cusp}}(\Gamma \backslash \text{SL}(2, \mathbb{R})))$ we may construct the L-series which is an element in $S_k(\Gamma)$; and conversely, starting from some a modular form $f \in S_k(\Gamma)$ we construct the corresponding L-series L_f .

There exists a unique intertwining operator A such that the L-series of which is equal to L_f .

3.1. Geometric Langlands Correspondence

The general Geometric Langlands Conjecture was stated by V. Drinfel'd and was then proven by E. Frenkel, D. Gaitsgory and Vilonen [FGV] and became the Geometric Langlands Correspondence (GLC). We will start to specify the general GLC in our particular case of group $\text{SL}_2(\mathbb{R})$.

Theorem 3.3 (Geometric Langlands Correspondence). There is a bijection

$$[\pi_1(\Sigma), \text{SO}(3)] \longleftrightarrow A(\text{SL}(2, \mathbb{R})) \quad (3.10)$$

between the set of equivalence classes of representation of the fundamental group $\pi_1(\Sigma)$ of the Riemannian surface $\Sigma = \Gamma \backslash \text{SL}(2, \mathbb{R})/\text{SO}(2)$ in $\text{SO}(3)$ and the set $A(\text{SL}(2, \mathbb{R}))$ of equivalence classes of automorphic representations of $\text{SL}(2, \mathbb{R})$

Proof: The theorem was proven in the general context of a reductive group in the works of [FGV]. The Geometric Langlands Correspondence is the one-to-one correspondence between the homotopy classes from the fundamental group $\pi_1(\Sigma)$ of the Riemannian surface Σ to the Langlands dual group ${}^L G$ and the set of equivalent classes of automorphic representations of the reductive group G . The machinery is very complicated involving the theory of Langlands dual groups, theory of auto-morphic forms on real reductive groups. For the particular case of $\text{SL}(2, \mathbb{R})$ many things are simplified, what we want to point out here

The main idea to prove the theorem is consisting of the following ingredients:

(1) It is well-known that every element of $SO(3)$ is conjugate with some element of the maximal torus subgroup $SO(2)$. Therefore the set of homomorphism from $\pi_1(\Sigma)$ into $SO(3)$ is the same as the set of character of the Borel subgroup (minimal parabolic subgroup) from which the discrete series rep-representations are induced.

Lemma 3.4. *There is one-to-one correspondence between the congruacy classes of $SO(3)$ in itself and the inducing character for the discrete series representations.*

(2) Moreover, the Γ invariance condition is the same as the condition to be extended from the local character of some automorphic component. The following two lemmas are more or less known [?].

Lemma 3.5. *Every representation of $\pi_1(\Sigma)$ in $SO(3)$ is defined by a system of conjugacy classes in $SO(3)$, one per generator of $\pi_1(\Sigma)$.*

Lemma 3.6. *Every system of conjugacy classes in the previous lemma 3.5 defines a unique modular form on H and hence a unique automorphic representation of $SL_2(\mathbb{R})$.*

3.2. Geometric Quantization

The idea of realizing the automorphic representations of reductive Lie groups was done in [D1]. In this section we show the concrete computation for the case of $SL(2, \mathbb{R})$

Theorem 3.7. The automorphic representations are obtained from the quantization procedure of fields based on geometric Langlands correspondence.

Proof: The discrete series representations can be realized through the geometric quantization as follows.

(1) The representation space of discrete series representation is consisting of square-integrable holomorphic functions

$$f(z) = \sum_{n=0}^{\infty} c_n z^n; \quad \sum_{n=0}^{\infty} |c_n|^2 < \infty \quad (3.11)$$

Lemma 3.8. *The Hardy space of holomorphic square-integrable functions can be realized as the exponential Fock space of the standard representation of $SO(2)$ in \mathbb{C} .*

Indeed every module-square convergence series of type 3.12 can be express as some element

$$f(z) = \sum_{n=0}^{\infty} n! c_n \frac{z^n}{n!}; \quad \sum_{n=0}^{\infty} |c_n|^2 < \infty \quad (3.12)$$

in the exponential vector space

$$EXPC = \sum_{n=0}^{\infty} \frac{C^{\otimes n}}{n!}, \quad C^{\otimes n} \cong C \quad (3.13)$$

(2) Let us now explain how can we obtain the representations as some results of geometric quantization procedure exposed in [D1]. The main idea of quantization of fields is the following: present the Lie algebra as some loop algebra over an appropriate Riemann surface by reduction, Kapustin-Witten and then combine the construction of the internal symmetry of by geometric quantization method with the construction of positive energy highest weight representation of loop algebra. Again in the paper, the general construction was explained, but we want to specify the deal in our concrete situation of $sl_2(\mathbb{R})$. The Lie algebra $sl_2(\mathbb{R})$ with 3 generators $sl_2(\mathbb{R}) = \langle H, X, Y \rangle_{\mathbb{R}}$ in the induced representations of discrete series act through the action of one-parameters subgroups.

$$g_3(t) = \exp(tH) = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix} \quad (3.14)$$

$$g_1(t) = \exp(t(X - Y)) = \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix}$$

$$g_2(t) = \exp(t(X + Y)) = \begin{pmatrix} \cosh t & \sinh t \\ \sinh t & \cosh t \end{pmatrix}$$

Under the representation π_n^\pm we have

$$\pi_n^\pm(g_k(t)) = e^{it\hat{U}_k} \tag{3.15}$$

and we have the relations

$$i\hat{U}_3 = \hat{H} = 2z \frac{\partial}{\partial z} + (n+1), \tag{3.16}$$

$$i\hat{U}_1 = \hat{X} - \hat{Y} = -(1+z^2) \frac{\partial}{\partial z} - (n+1)z,$$

$$i\hat{U}_2 = \hat{X} + \hat{Y} = (1-z^2) \frac{\partial}{\partial z} - (n+1)z.$$

In this representation, the action of the element

$$\Delta = \frac{-1}{4}(\hat{U}_1^2 - \hat{U}_2^2 - \hat{U}_3^2) = \tag{3.17}$$

$$= \frac{-1}{4} \left((\hat{X} - \hat{Y})^2 - (\hat{X} + \hat{Y})^2 - (\hat{H})^2 \right) = \frac{1}{4}(\hat{H}^2 + 4\hat{X}\hat{Y}) =$$

$$= (z - \bar{z})^2 \frac{\partial^2}{\partial z \partial \bar{z}} = -y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)$$

This action can be represented as the action of the loop algebra over the Riemann surface Σ with values in $SO(2)$, the elements of the loop algebra are presented in form of some formal/conformal Laurent series with values in the corresponding Lie algebra of form of connection appeared in the construction of induced representations, i.e.

$$T(z) = \sum_{n=-\infty}^{\infty} c_n z^n, c_n \in so(2), z \in \Sigma. \tag{3.18}$$

In our case the Lie algebra $so(2)$ is one dimensional and we have all number coefficients $c_n, n \in \mathbb{Z}$.

Lemma 3.9. *The decomposition 3.13 of $EXP \mathbb{C}$ presents the weight decomposition of $sl_2(\mathbb{R})$ in which H keeps each component, X acting as creating operator and Y is acting as some annihilating operator.*

(3) The lowest weight representations of the loop algebras are realized through the lowest weight representations of the Virassoro algebra as follows. Let us consider the generators.

$$L_n = \int_{\Sigma} z^{n+1} T(z) dz \tag{3.19}$$

for any element

$$T(\cdot) : \Sigma \rightarrow SO(2)$$

from the loop algebra presentation are realized in the Fock space of the standard representation. These generators satisfy the Virassoro algebra relations.

$$[L_m, L_n] = (n-m)L_{m+n} + \delta_{n,-m} \frac{n(n^2-1)}{12} L_0, \tag{3.20}$$

where in an irreducible representation, $Z = cL_0 = cI$ is the central charge element.

The proof of Theorem 3.7 is therefore achieved.

IV. ARTHUR-SELBERG TRACE FORMULA

4.1. Trace Formula

Let us review the trace formula due to Selberg and J. Arthur. Remind that by $H = SL(2, \mathbb{R}) \backslash SO(2)$ denote the upper Poincaré half-plane,

$$\mathbb{H} = \{z = x + iy \in \mathbb{C} \mid \Im z = y > 0\},$$

Γ denote a Langlands type discrete subgroup of finite type with finite number of cusps $\kappa_1, \dots, \kappa_h$. Let

$$\Gamma_0 = \left\{ \gamma = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \mid b \in \mathbb{Z} \right\} \subset SL(2, \mathbb{Z})$$

And

$$\Gamma_i = \{ \sigma \in \Gamma \mid \sigma \kappa_i = \kappa_i \} \subset SL(2, \mathbb{Z})$$

and $\sigma_i \in SL(2, \mathbb{R})$, $i = \overline{1, h}$ are such that $\sigma_i \Gamma_i \sigma_i^{-1} = \Gamma_0$. $\sigma_i \Gamma_i \sigma_i = \Gamma_0$, $\mathfrak{S} = L^2(\Gamma \backslash G)$ denote the space of square-integrable functions on G on which there is a natural regular representation R of G by formula.

$$[R\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right)f](z) = f\left(\frac{az + b}{cz + d}\right), z \in \mathbb{H}.$$

In particular, the unipotent radical of $N = \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mid x \in \mathbb{R} \right\}$ is acting by the translation on variable z to $z + x$. For any function $\psi: \mathbb{N} \backslash \mathbb{H} \rightarrow \mathbb{C}$ of variable y decreasing fast enough when y approaches to 0 or ∞ , one defines the incomplete θ -series.

$$\theta_{t, \psi}(z) = \sum_{\sigma \in \Gamma_i \backslash \Gamma} \psi(\sigma_i^{-1} \sigma z)$$

which is certainly of class $L^2(N \cap \Gamma \backslash H)$. Denote by

$$\Theta = \text{h}\theta_{t, \psi} \mid \forall \psi, t_i \in L^2(\Gamma \backslash H)$$

the space of incomplete θ -series.

It is well-known that the orthogonal complement Θ^\perp of Θ in $L^2(\Gamma \backslash H)$ is isomorphic to the space H_0 of cuspidal parabolic forms or in other words, of automorphic forms with zero Fourier constant terms of automorphic representations, $H = \Theta \oplus H_0$. Consider the Hecke algebra $\mathcal{H}(SL(2, \mathbb{R}))$ of all convolution Hecke operators of form as follows. Let $F: \mathbb{H} \rightarrow \mathbb{C}$ be a function K invariant with respect to transformations of form $z \rightarrow \gamma z$, for all $\gamma \in K$. Such a function is uniquely defined by a function on $K \backslash G / K$, or a so called spherical function F on G which is left and right invariant by K . Under convolution these functions provide the Hecke algebra of Hecke operators by convolution with functions in representations. All the Hecke operators have kernel as follows. For any automorphic function f .

$$f(\gamma z) = f(z), \forall \gamma \in \Gamma.$$

$$\begin{aligned} (F * f)(z) &= \int_{\mathbb{H}} F(g'^{-1}g)f(g')dg' = \int_{\mathbb{H}} k(z, z')f(z')dz' = \int_{\Gamma \backslash \mathbb{H}} \sum_{\gamma \in \Gamma} k(z, \gamma z')f(z')dz', \\ &= \int_{\Gamma \backslash \mathbb{H}} K(z, z')f(z')dz', \end{aligned}$$

where $z = gi, z'g'i, i = \sqrt{-1}$,

$$K(z, z') = \sum_{\gamma \in \Gamma} k(z, \gamma z').$$

Denote the sum of kernel over cups by

$$H(z, z') = \sum_{i=1}^h H_i(z, z') = \sum_{i=1}^h \sum_{\gamma_i \in \Gamma_i \backslash \Gamma} \int_{-\infty}^{+\infty} K(z, \gamma_i n(x) \gamma_i^{-1} \gamma z') dx$$

The Hecke operator with kernel $K(z, z')$ has the same spectrum as the operator with kernel $K^*(z, z') = K(z, z') - H(z, z')$. The kernel $K^*(z, z')$ are bounded and the fundamental domain $\Gamma \backslash \mathbb{H}$ has finite volume, therefore the kernels $K^*(z, z')$ are of class L^2 on $D \times D$, $D = \Gamma \backslash \mathbb{H}$, and all the Hecke operators are compact operators. The Hecke operators keeps each irreducible components of Θ^\perp invariant and therefore are scalar on each automorphic representation. On each irreducible component, the Laplace operators has also a fixed eigenvalue.

$$\Delta f = \lambda f, \lambda = \frac{s(s-1)}{4}, \Delta = -y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)$$

One deduces therefore the theorem of spectral decomposition for the discrete part of the regular representation.

Theorem 4.1 (Spectral decomposition). In the induced representation space of $\text{Ind}_B^G \chi, \lambda, \varepsilon$

, choose

$$H_n = \left\{ f \in H \mid \pi \left(\begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \right) f = e^{in\theta} f \right\}.$$

They are all of dimension 1 and

$$H = \bigoplus_{n \in \mathbb{Z}} H_n.$$

The discrete part $\text{RL}_{\text{cusp}}^2(\Gamma \backslash \text{SL}(2, \mathbb{R}))$ of the regular representation can be decomposed as the sum of the discrete series representations π_n^\pm in spaces

$$D_{s+1}^+ = \bigoplus_{n \equiv \varepsilon \pmod{2}, n \geq m} H_n$$

or

$$D_{s+1}^- = \bigoplus_{n \equiv \varepsilon \pmod{2}, n \leq -m} H_n,$$

$s \in \mathbb{Z}$, $s > 0$ and $s+1 \equiv \varepsilon \pmod{2}$ and there exists $m \in \mathbb{Z}$, $m = s+1$, $m > 0$, induced from $\chi^{\lambda, \varepsilon} = |a|^{\lambda} (\text{sign } a)^\varepsilon$ and limits of principal series representations π_0^\pm in D_1^+ or D_1^- as two component of the representation $\pi_1^0 = \text{Ind}_B^G \chi, \lambda, \varepsilon$, induced from the character $\chi_{0,1}$.

Remark 4.2. In the spaces $\bigoplus_{-m < n < m} H_n$ of dimension $2m - 1$ the finite dimensional representations V_m are realized.

Corollary 4.3. For any function ϕ of class $C_c^\infty(G)$, the operator $\pi_n^\pm(\phi)$ is of trace class and is a distribution denoted by θ_n^\pm (following Harish-Chandra) which are uniquely defined by their restriction to the maximal compact subgroup $K = \text{SO}(2)$ and

$$\text{tr } R(\varphi) = \sum_{n \in \mathbb{Z}, n \geq 0, \pm} m(\pi_n^\pm) \theta_n^\pm(\varphi),$$

with multiplicities $m(\pi_n^\pm)$

4.2. Stable Trace Formula

The Galois group $\text{Gal}(\mathbb{C}/\mathbb{R}) = \mathbb{Z}_2$ of the complex field \mathbb{C} is acting on the discrete series representation by character $\kappa(\sigma) = \pm 1$. Therefore the sum of characters can be rewrite as some sum over stable classes of characters.

$$\text{tr } R(f) = \sum_{n=1}^{\infty} \sum_{\epsilon=\pm 1} (\Theta_n^+(f) - \Theta_n^-(f)).$$

Remark 4.4. The stable trace formula is uniquely defined by its restriction to the maximal compact subgroup $K = \text{SO}(2)$ and is

$$S\Theta_n = \frac{\Theta_n^+ - \Theta_n^-}{e^{i\theta} - e^{-i\theta}} - 2i \sin \theta (e^{in\theta} - e^{-in\theta}).$$

V. ENDOSCOPYS

5.1. Stable Orbital Integral

Let us remind that the orbital integral is defined as

$$\mathcal{O}_\gamma(f) = \int_{G_\gamma \backslash G} f(x^{-1}\gamma x) dx$$

The main idea of computation of endoscopy transfer was explained in [La], we make it here in more detail to clarify some points.

The complex Weyl group is isomorphic to \mathfrak{S}_2 while the real Weyl group is iso-morphic to \mathfrak{S}_1 . The set of conjugacy classes inside a strongly regular stable elliptic conjugacy class is in bijection with the pointed set $\mathfrak{S}_2 / \mathfrak{S}_1 = \mathfrak{S}_2$ that can be viewed as a sub-pointed-set of the group $\mathfrak{C}(\mathbb{R}, T, G) = (Z_2)$ we shall denote by $\mathfrak{K}(\mathbb{R}, T, G) \cong \mathbb{Z}_2$ its Pontryagin dual.

Consider $\kappa \neq 1$ in $\mathfrak{K}(\mathbb{R}, T, G)$ such that $\mathcal{K}(H) = -1$. Such a κ is unique. The endoscopic group H one associates to κ is isomorphic to $\text{SO}(2)$ the positive root of \mathfrak{h} in H (for a compatible order) being $\alpha = \rho$.

Let f_μ be a pseudo-coefficient for the discrete series representation π_μ then the κ -orbital integral of a regular element γ in $T(\mathbb{R})$ is given by

$$\mathcal{O}_\gamma^\kappa(f_\mu) = \int_{G_\gamma \backslash G} \kappa(x) f_\mu(x^{-1}\gamma x) dx = \sum_{\text{sign}(w)=1} \kappa(w) \Theta_\mu^G(\gamma_w^{-1}) = \sum_{\text{sign}(w)=1} \kappa(w) \Theta_{w\mu}(\gamma^{-1}),$$

because there is a natural bijection between the left coset classes and the right coset classes.

5.2. Endoscopic Transfer

We make details for the guides of J.-P. Labesse [La]. The simplest case is the case when $\gamma = \text{diag}(a, a^{-1})$. In this case, because of Iwasawa decomposition $x = auk$, and the K -bivariance, the orbital integral is

$$\begin{aligned} \mathcal{O}_\gamma(f) &= \int_{G_\gamma \backslash G} f(x^{-1}\gamma x) dx = \int_U f(u^{-1}\gamma u) du \\ &= \int_{\mathbb{R}} f\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right) dx \\ &= \int_{\mathbb{R}} f\left(\begin{pmatrix} 1 & -x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right) dx \\ &= \int_{\mathbb{R}} f\left(\begin{pmatrix} a & (a - a^{-1})x \\ 0 & a^{-1} \end{pmatrix}\right) dx = |a - a^{-1}|^{-1} \mathcal{O}_\gamma(\tilde{f}), \end{aligned}$$

where \tilde{f} is the result of integration on variable x . The integral is absolutely and uniformly convergent and therefore is smooth function of $a \in \mathbb{R}_+^*$. Therefore the function.

$$f^H(\gamma) = \Delta(\gamma)\mathcal{O}_\gamma(f), \quad \Delta(\gamma) = |a - a^{-1}|$$

is a smooth function on the endoscopic group $H = \mathbb{R}^*$

The second case is the case where $\gamma = k_\theta \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix}$. We have again, $x = auk$ and

$$\begin{aligned} \mathcal{O}_{k(\theta)}(f) &= \int_{G_{k(\theta)} \backslash G} f(k^{-1}u^{-1}a^{-1}k(\theta)auk)dx \frac{dy}{|y|}d\theta \\ &= \int_{G_{k(\theta)} \backslash G} f(u^{-1}a^{-1}k(\theta)au)dx \frac{dy}{|y|}d\theta \\ &= \int_{G_{k(\theta)} \backslash G} f\left(\begin{pmatrix} 1 & -x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} a^{-1} & 0 \\ 0 & a \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right)dx \frac{dy}{|y|}d\theta \\ &= \int_1^\infty \bar{f}\left(\begin{pmatrix} \cos\theta & t \sin\theta \\ -t^{-1} \sin\theta & \cos\theta \end{pmatrix}\right)dt, \end{aligned}$$

\bar{f} is the result of integration on variable x . When f is an element of the Hecke algebra, i.e. f is of class $C_0^\infty(G)$ and is K -bi-invariant, the integral is converging absolutely and uniformly. Therefore the result is a function $F(\sin\theta)$. The function f has compact support, then the integral is well convergent at $+\infty$. At the another point 0, we developpe the function F into the Taylor-Lagrange of the first order with respect to $\lambda = \sin\theta \rightarrow 0$

$$F(\lambda) = A(\lambda) + \lambda B(\lambda),$$

where $A(\lambda) = F(0)$ and $B(\lambda)$ is the error-correction term $F'(\tau)$ at some intermediate value τ , $0 \leq \tau \leq t$. Remark that

$$\begin{pmatrix} \sqrt{1-\lambda^2} & t\lambda \\ -t^{-1}\lambda & \sqrt{1-\lambda^2} \end{pmatrix} = \begin{pmatrix} t^{1/2} & 0 \\ 0 & t^{-1/2} \end{pmatrix} \begin{pmatrix} \sqrt{1-\lambda^2} & \lambda \\ \lambda & \sqrt{1-\lambda^2} \end{pmatrix} \begin{pmatrix} t^{-1/2} & 0 \\ 0 & t^{1/2} \end{pmatrix}$$

we have

$$\begin{aligned} B &= \frac{dF(\tau)}{d\lambda} = \frac{d}{d\lambda} \int_0^{+\infty} \text{sign}(t-1)f\left(\begin{pmatrix} \sqrt{1-\lambda^2} & t\lambda \\ -t^{-1}\lambda & \sqrt{1-\lambda^2} \end{pmatrix}\right)dt \Big|_{t=\tau} \\ &= \int_0^{+\infty} \text{sign}(t-1)g\left(\begin{pmatrix} \sqrt{1-\lambda^2} & t\lambda \\ -t^{-1}\lambda & \sqrt{1-\lambda^2} \end{pmatrix}\right)\frac{dt}{t}, \end{aligned}$$

where $g \in C_c^\infty(N)$ and $g(\lambda) \sim O(-t^{-1}\lambda)^{-1}$. B is of logarithmic growth and

$$B(\lambda) \cong \ln(|\lambda|^{-1})g(1)$$

up to constant term, and therefore is continuous.

$$A = F(0) = |\lambda|^{-1} \int_0^\infty f\left(\begin{pmatrix} 1 & \text{sign}(\lambda)u \\ 0 & 1 \end{pmatrix}\right)du - 2f(I_2) + o(\lambda)$$

Hence the functions

$$G(\lambda) = |\lambda|(F(\lambda) + F(-\lambda)),$$

$$H(\lambda) = \lambda(F(\lambda) - F(-\lambda))$$

have the Fourier decomposition

$$G(\lambda) = \sum_{n=0}^N (a_n |\lambda|^{-1} + b_n) \lambda^{2n} + o(\lambda^{2N})$$

$$H(\lambda) = \sum_{n=0}^N h_n \lambda^{2n} + o(\lambda^{2N})$$

Summarizing the discussion, we have that in the case of $\gamma = k(\theta)$, there exists also a continuous function f^H such that

$$f^H(\gamma) = \Delta(\gamma)(O\gamma(f) - O_w\gamma(f)) = \Delta(k(\theta))SO\gamma(f),$$

where $\Delta(k(\theta)) = -2i \sin \theta$.

Theorem 5.1. *There is a natural function $\varepsilon : \Pi \rightarrow \pm 1$ such that in the Grothendieck group of discrete series representation ring,*

$$\sigma_G = \sum_{\pi \in \Pi} \varepsilon(\pi) \pi,$$

the map $\sigma \rightarrow \sigma_G$ is dual to the map of geometric transfer, that for any f on G , there is a unique f^H on H

$$\text{tr } \sigma_G(f) = \text{tr } \sigma(f^H).$$

Proof.: There is a natural bijection $\Pi_\mu \cong \mathfrak{D}(\mathbb{R}, H, G)$, we get a pairing

$$\langle \cdot, \cdot \rangle : \Pi_\mu \times k(R, H, G) \rightarrow \mathbb{C}.$$

Therefore we have

$$\text{tr}_{\Sigma_\nu}(f^H) = \sum_{\pi \in \Pi_\Sigma} \langle s, \pi \rangle \text{tr } \pi(f).$$

Suppose given a complete set of endoscopic groups $H = \mathbb{S}^1 \times \mathbb{S}^1 \times \{\pm 1\}$ or $SL(2, \mathbb{R}) \times \{\pm 1\}$. For each group, there is a natural inclusion

$$\eta: {}^L H \hookrightarrow {}^L G$$

Let $\phi : DW_{\mathbb{R}} \rightarrow {}^L G$ be the Langlands parameter, i.e. a homomorphism from the Weil-Deligne group $DW_{\mathbb{R}} = W_{\mathbb{R}} \rtimes \mathbb{R}_+^*$ the Langlands dual group, $S\phi$ be the set of conjugacy classes of Langlands parameters modulo the connected component of identity map. For any $s \in S\phi$, $\tilde{H}s = \text{Cent}(s, \tilde{G})^\circ$ the connected component of the centralizer of $s \in S\phi$ we have $\tilde{H}s$ is conjugate with H . Following the D. Shelstad pairing.

$$\begin{aligned} \langle s, \pi \rangle : S_\phi \times \Pi(\phi) &\rightarrow \mathbb{C} \\ \varepsilon(\pi) &= c(s) \langle s, \pi \rangle. \end{aligned}$$

Therefore, the relation

$$\sum_{\sigma \in \Sigma_s} \text{tr } \sigma(f^H) = \sum_{\pi \in \Pi} \varepsilon(\pi) \text{tr } \pi(f)$$

can be rewritten as

$$\tilde{\Sigma}_s(f^H) := \sum_{s \in \Pi} \langle s, \pi \rangle \text{tr } \pi(f)$$

and

$$\tilde{\Sigma}_s(f^H) := c(s)^{-1} \sum_{\sigma \in \tilde{\Sigma}_s} \text{tr } \sigma(f^H).$$

We arrive, finally to the result

Theorem 5.2.

$$\text{tr } \pi(f) = \frac{1}{\#S} \sum_{\varphi} \langle s, \pi \rangle_{\Sigma_s}(\hat{f}^H).$$

VI. POISSON SUMMATION FORMULA

In the Langlands picture of the trace formula, the trace of the restriction of the regular representation on the cuspidal parabolic part is the coincidence of the spectral side and the geometric side. We refer the reader to the work of J.-p.

Labesse [La] for more detailed exposition

$$\sum_{\pi} m(\pi) \hat{f}(\pi) = \sum_{\gamma \in \Gamma \cap H} a_{\gamma} \hat{f}(\gamma) \tag{6.1}$$

Let us do this in more details.

6.1. Geometric Side of the Trace Formula

Theorem 6.1. The trace formula for the regular representation of $SL(2, \mathbb{R})$ in the space of cusp forms is decomposed into the sum of traces of automorphic representations with finite multiplicities is transferred into the modified Poisson summation formula.

$$\sum_{\gamma \in \Gamma \cap H} \varepsilon(\gamma) \mathcal{O}_{\gamma}(f) = \sum_{\gamma \in \Gamma \cap H} \varepsilon(\gamma) \text{vol}(\Gamma \cap H) \int_{H \backslash G} f(x^{-1} \gamma x) dx \tag{6.2}$$

Proof. It is easy to see that the restriction of the Laplace operator Δ on the Cartan subgroup H is elliptic and therefore the Cauchy problem for the other variables has a unique solution. The solution is the trace formula for the cuspidal parabolic part of the regular representation.

$$\text{tr } R(f)|_{L^2_{\text{cusp}}(\Gamma \backslash G)} = \sum_{\gamma \in \Gamma \cap H} \varepsilon(\gamma) \mathcal{O}_{\gamma}(f) \tag{6.3}$$

From another side we have

$$\begin{aligned} \text{tr } R(f)|_{L^2_{\text{cusp}}(\Gamma \backslash G)} &= \sum_{\gamma \in \Gamma \cap H} \varepsilon(\gamma) \text{vol}(\Gamma \cap H) \int_{H \backslash G} f(x^{-1} \gamma x) dx \\ &= \sum_{\gamma \in \Gamma \cap H} \varepsilon(\gamma) \text{vol}(\Gamma \cap H) \mathcal{S} \mathcal{O}_{\gamma}(f^H). \end{aligned} \tag{6.4}$$

6.2. Spectral Side Of The Trace Formula

The following result is well-known, see e.g. ([GGPS], Ch. 1).

Theorem 6.2 (Gelfand - Graev - Piatetski-Shapiro). For any compactly supported function $f \in C_0^{\infty}(SL_2(\mathbb{R}))$ the operator $R(f)|_{L^2_{\text{cusp}}(\Gamma \backslash SL_2(\mathbb{R}))}$ is of trace class and each irreducible component is of finite multiplicity.

$$R(f)|_{L^2_{\text{cusp}}(\Gamma \backslash SL_2(\mathbb{R}))} = \sum_{\pi \in \mathcal{A}(SL_2(\mathbb{R}))} m(\pi) \pi(f), \tag{6.5}$$

where $m(\pi) = \dim_{\mathbb{C}} \text{Hom}_{SL_2(\mathbb{R})}(D_k, L^2_{\text{cusp}}(\Gamma \backslash SL_2(\mathbb{R})))$.

6.3. Poisson Summation Formula

Therefore, following the Poisson summation formula we have the equality of the both sides. Let us denote by χ_k the character of $SO(2)$ that induce the discrete series representation D_k of $SL_2(\mathbb{R})$. In our case the Cartan subgroup is $SO(2)$ and we have the ordinary.

Lemma 6.3. De note the universal covering of $H = SO(2)$ by $\tilde{H} = Spin(2)$.

$$\sum_{\gamma \in \Gamma \cap \tilde{H}} \delta(x + \gamma) = \sum_{(\varepsilon, k) \in \mathbb{Z}_2 \times \mathbb{Z} = \hat{H}} (\text{sign } x)^\varepsilon \chi_k(x) \quad (6.6)$$

Proof. It is the same as ordinary Poisson summation, lifted to the universal covering \tilde{H} of $SO(2)$.

Theorem 6.4. The trace $\text{tr } R(f)$ of the restriction of the regular representation on the cuspidal parabolic part $L^2_{\text{cusp}}(G)$ is computed by the formulas

$$\sum_{\pi \in \mathcal{A}(G)} m(\pi) \pi(f) = \sum_{\substack{\gamma \in \Gamma \cap \tilde{H} \\ \varepsilon(\gamma) = \pm 1}} \varepsilon(\gamma) \text{vol}(\Gamma \cap \tilde{H}) \mathcal{O}_\gamma(f) = \sum_{\gamma \in \Gamma \cap H} \text{vol}(\Gamma \cap H) S \mathcal{O}_\gamma(f^H) \quad (6.7)$$

Proof. The proof just is a combination of the previous theorems 6.1, 6.2 and 6.3 therefore is complete.

VII. CONCLUSION

We developed in this paper a new approach to the construction of automorphic representations based on a procedure of quantization. The Langlands picture of automorphic representations and endoscopy are precised for the case of $SL(2, \mathbb{R})$.

REFERENCES

- [1] [B] A. Borel, Automorphic forms on $SL_2(\mathbb{R})$, Cambridge Univ. Press, Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore, Sao Paulo, 2008.
- [2] [D1] Do Ngoc Diep, A Quantization Procedure of fields based on Geometric Langlands Correspondence, Intl. J. Math. Mathl. Sci., 2009, 14 pages, doi:10.1155/2009/749631.
- [3] [FGV] E. Frenkel, D. Gaitsgory, D. Kazhdan, and K. Vilonen, Geometric realization of Whittaker functions and the Langlands Conjecture, J. Amer. Math. Soc. 11(2), 1998, 451-484.
- [4] [GGPS] I. Gelfand, M. Graev, Y. Piatetski-Shapiro, Representation Theory and Automorphic Functions, Generalized Functions Series, Vol 6, Nauka Press, Moscow, 1969.
- [5] [La] J.-P. Labesse, Introduction to Endoscopy, Snowbird Lectures, June 2006 www.institut.math.jussieu.fr/projets/fa/bpFiles/Labesse.pdf.
- [6] [L] S. Lang, $SL_2(\mathbb{R})$, Springer-Verlag New York Berlin Heidelberg Tokyo, Addison-Wesley, 1975