American Research Journal of
Sports Medicine

ARJ PUBLISHING
Inspiring Research. Shaping Futures

Review Article

Neuromotor Adaptations and Functional Athletic Capacity in Professional Football: An Integrated Approach to Performance Restoration

Hartmut Frosh[1*], Alexander Behr[2]

[1]Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK.
[2]Sports Medicine Ultrasound Group (SMUG), London, UK.

Correspondence to: Hartmut Frosh, Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK.

Publication Information: Accepted: May 19, 2026**, Published:** May 26, 2026

Citation: Frosh H, et al. Neuromotor Adaptations and Functional Athletic Capacity in Professional Football: An Integrated Approach to Performance Restoration. American Research Journal of Sports Medicine, Vol 1, no. 1, 2026: 1-7.

Copyright © 2026 Frosh H, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Modern professional football places substantial demands on athletes' physical capacity, perceptual processing, movement coordination, and rapid decision-making. The interaction between these abilities becomes especially important during competitive match situations and the rehabilitation of players recovering from major musculoskeletal injuries. Conventional rehabilitation strategies frequently emphasize muscular strength, joint stability, and running performance, while cognitive and sensorimotor adaptations may receive less attention. This review examines the contribution of visual processing, attentional regulation, neuromuscular conditioning, blood flow restriction (BFR), ischemic preconditioning (IPC), and virtual reality (VR) to athletic development and injury rehabilitation. emphasis is placed on divided attention, selective information processing, anticipatory responses, and movement regulation under unpredictable conditions. The potential combination of BFR and VR is also discussed as an emerging rehabilitation strategy when conventional physical loading is restricted. Integrating perceptual-motor activities with progressive conditioning may improve functional preparedness and support a more comprehensive return-to-sport process. Further controlled investigations are required to establish the effectiveness, safety, and practical application of combined interventions in professional football.

Keywords: Blood Flow Restriction; Cognitive-Motor Integration; Elite Football; Ischemic Preconditioning; Neuromuscular Rehabilitation; Return To Play; Virtual Reality; Visual Perception.

INTRODUCTION

Professional football has undergone considerable changes in physical intensity, tactical organization, and competitive expectations. Players are increasingly required to perform repeated high-speed movements, accelerate rapidly, change direction, and maintain technical accuracy while responding to continuously changing match situations. These requirements demonstrate the relationship between physical conditioning and the perceptual abilities necessary for effective sporting performance. Match transitions, pressing sequences, and rapid counterattacking movements create additional physiological and cognitive demands that influence individual and collective performance [1].

The increasing intensity of competitive football has also changed the physical requirements associated with different playing positions. High-speed running, acceleration, deceleration, and repeated high-intensity actions contribute to the workload experienced during competition. Consequently, conditioning programs must address the specific physiological requirements of individual positions rather than relying exclusively on generalized fitness development [2].

Injury rehabilitation presents additional challenges because restoring muscular strength and joint function does not necessarily indicate complete recovery of movement regulation or cognitive performance. Anterior

cruciate ligament (ACL) injury, for example, can produce changes in sensory information processing, cortical activity, and neuromuscular coordination. These adaptations may influence movement execution even after athletes regain satisfactory physical capacity [3]. Neuroplastic changes following injury can affect the interaction between the central nervous system and the musculoskeletal system. Rehabilitation programs that incorporate motor learning, progressive movement challenges, and cognitive engagement may therefore provide a broader approach to restoring functional performance [4].

Contemporary rehabilitation frameworks increasingly recognize the importance of combining physical conditioning with perceptual-motor development. This approach supports the progressive introduction of attention-demanding activities, decision-making tasks, and unpredictable movement situations during recovery [5].

LITERATURE REVIEW

Cognitive and Physical Requirements of Competitive Football

Football performance depends on the coordinated functioning of physical, technical, tactical, and perceptual abilities. Players must continuously interpret environmental information, recognize developing opportunities, anticipate opponents' actions, and select appropriate movement responses.

Working memory contributes to the temporary storage and manipulation of information required during complex sporting activities. It enables players to retain tactical instructions, monitor surrounding movements, and adjust their actions according to changing match conditions [6]. Selective attention supports the identification of relevant information while reducing interference from unnecessary environmental stimuli. In football, this process enables athletes to recognize passing opportunities, identify defensive threats, and respond efficiently to changes in possession [7].

The interaction between cognitive processing and muscular function is also important following ACL injury. Alterations in cortical control and quadriceps activation have been reported in individuals with previous knee injuries, indicating that rehabilitation should address neurological and muscular components together [8].

Visual Processing and Perceptual-Motor Development

Visual information provides an essential foundation for movement preparation and decision-making in football. Players continuously interpret the position of teammates, opponents, and the ball before selecting an appropriate technical or tactical response. Virtual and augmented reality applications have attracted increasing interest in sports rehabilitation because they allow athletes to practice perception, movement planning, and decision-making within controlled environments. These technologies may also provide alternative training opportunities when conventional field-based activities are temporarily restricted [9].

Positional requirements further influence the type of information athletes must process. Midfielders frequently monitor several passing options, defenders evaluate opponent movements, and forwards anticipate spaces behind defensive lines. These differences support the inclusion of position-specific perceptual activities within physical preparation programs [10].

Blood Flow Restriction and Neuromuscular Conditioning

Blood flow restriction training involves applying external pressure to a limb to partially limit arterial inflow and restrict venous return during exercise. This method allows athletes to undertake low-load resistance activities while generating substantial metabolic stress.

Vascular occlusion produces changes in muscle oxygenation and local physiological conditions, which contribute to the responses associated with restricted-flow exercise [11].

Research involving ischemia-reperfusion interventions has examined their potential influence on exercise performance and post-exercise recovery. These findings have encouraged further investigation into their application within athletic conditioning [12]. Low-intensity resistance exercise combined with BFR has demonstrated potential benefits for aerobic conditioning and physical performance. This approach may be useful when conventional high-intensity loading cannot be tolerated [13].

Muscular adaptations associated with heavy resistance exercise and BFR-based training include changes in muscle size and functional capacity. The magnitude of these responses depends on training intensity, intervention duration, and individual characteristics [14]. Appropriate application requires careful control of cuff pressure, exercise volume, recovery intervals, and participant suitability. Standardized procedures are necessary to improve consistency and minimize potential complications [15].

 © 2026 Frosh H

Ischemic Preconditioning and Athletic Performance

Ischemic preconditioning involves repeated periods of temporary circulatory restriction followed by reperfusion. It has been investigated as a method of influencing physiological responses before subsequent physical activity. Research involving older adults has examined the relationship between restricted-flow exercise, cognitive performance, and muscular function. These findings provide preliminary evidence for investigating combined physical and cognitive interventions, although their direct application to elite football remains uncertain [16]. The feasibility of garment-integrated BFR equipment has also been investigated, providing information about its practical use during lower-limb exercise. Such equipment may offer opportunities for controlled application in rehabilitation environments [17].

Low-load BFR cycling has demonstrated potential adaptations in muscle mass, strength, and aerobic capacity, supporting its investigation as an alternative conditioning strategy when high mechanical loading is unsuitable [18].

Physiological Mechanisms Associated with BFR

The physiological responses to BFR involve metabolic stress, altered oxygen availability, cellular swelling, and changes in local muscle activity. These mechanisms may contribute to adaptations associated with low-load resistance exercise. Research has also investigated the relationship between BFR and neurotrophic responses. Increased circulating brain-derived neurotrophic factor following lactate-related interventions provides a possible physiological connection between metabolic stimulation and neurobiological processes [19].

BFR-based walking interventions have also been investigated in populations with reduced physical function. These findings support further exploration of restricted-flow exercise as a method of maintaining physical activity when conventional exercise intensity is difficult to achieve [20].

However, evidence obtained from older or recreationally active populations should not automatically be generalized to professional footballers. Differences in training history, physical capacity, injury characteristics, and competitive workload must be considered.

Virtual Reality in Sports Rehabilitation

Virtual reality provides an interactive environment in which visual information and movement-related responses can be practiced through computer-generated simulations. Head-mounted displays and motion-tracking technologies allow users to interact with virtual objects and respond to changing visual stimuli. The combination of VR-based rehabilitation and BFR has been proposed as an approach for maintaining cognitive engagement while limiting mechanical loading. This strategy may be useful during periods when conventional exercise intensity must remain restricted [21].

Ischemic preconditioning has also been investigated as a method of improving maximal exercise performance, indicating its potential relevance to physical preparation and recovery [22]. Additional research has examined its application during resistance training, where improvements in exercise-session performance have been reported. These findings support further investigation into the relationship between circulatory preconditioning and muscular performance [23]. Repeated-sprint cycling research has also examined the influence of ischemic preconditioning on high-intensity exercise capacity. Such findings may provide a physiological basis for investigating its application in intermittent sports [24].

Cognitive-Motor Training and Return-to-Play Preparation

Virtual environments provide opportunities to examine sporting decisions under controlled conditions. They can reproduce elements of passing, ball movement, opponent positioning, and tactical responses while allowing researchers to regulate the complexity of visual information. A review of virtual reality applications in team-ball sports identified their potential for evaluating and developing sport-specific performance abilities [25]. The use of animated and immersive virtual environments has also been examined for assessing decision-making in team sports. Differences in simulation design and environmental realism may influence how effectively laboratory-based performance reflects actual match situations [26].

Virtual environments have additionally been investigated for evaluating functional walking difficulties. These applications demonstrate how controlled simulations can support movement assessment without requiring athletes to complete every task in a conventional physical environment. Research comparing personal-space behavior during obstacle avoidance in physical and virtual environments provides further insight into movement regulation under different environmental conditions.

 © 2026 Frosh H

Development of Attentional and Perceptual Abilities

Training activities should progressively challenge athletes to identify relevant information and select appropriate responses. Coaches can introduce visual signals, changing movement directions, and opponent-related stimuli into technical exercises. Divided-attention activities can combine ball control with external decision-making requirements. For example, a player may perform a passing sequence while responding to changing visual signals provided by the coach. Selective-attention exercises can require athletes to identify specific teammates, recognize defensive movements, or respond only to relevant visual cues. These activities should progress from predictable situations toward more complex exercises that reproduce the demands of competitive football (Table 1).

Table 1: Cognitive Processes and Their Application During Football Performance.

Cognitive component	Functional description	Practical football application
Anticipatory processing	Predicting an upcoming action from available information	Moving into space before a teammate delivers a pass
Selective attention	Identifying relevant information while disregarding distractions	Recognizing an opponent's movement during defensive play
Divided attention	Processing multiple information sources during activity	Controlling the ball while monitoring teammates
Visual scanning	Collecting information from different areas of the playing environment	Checking surrounding spaces before receiving possession
Response inhibition	Suppressing an unsuitable or premature action	Avoiding a pass when the intended route becomes blocked
Working memory	Retaining and manipulating information temporarily	Remembering tactical instructions while responding to opposition movements

Attacking and Defensive Training Situations

Attacking players must recognize available spaces, identify passing opportunities, and anticipate the movement of teammates and opponents. These abilities can be developed through small-sided games, positional exercises, and transition-based activities. Defensive players require efficient visual scanning and rapid responses to opponent movements. Exercises involving interception, tracking, pressing, and closing down spaces can reproduce these requirements (Table 2).

Table 2: Match-Specific Perceptual Activities for Attacking and Defensive Players.

Playing role	Training situation	Intended development
Central midfielder	Receiving the ball under pressure and identifying alternative passing routes	Visual scanning and decision-making
Striker	Anticipating crosses and moving into goal-scoring positions	Spatial awareness and anticipatory movement
Winger	Recognizing defensive gaps during counterattacks	Perceptual speed and movement selection
Central defender	Intercepting passes and responding to attacking movements	Anticipation and reaction control
Full-back	Tracking overlapping opponents	Visual monitoring and directional adjustment
Goalkeeper	Responding to changing shooting angles	Visual processing and rapid motor responses

Progressive Cognitive-Motor Rehabilitation

Rehabilitation following ACL reconstruction should progress according to the athlete's physical condition, movement quality, and functional readiness. Early rehabilitation generally emphasizes joint mobility, muscular activation, and basic movement control. As recovery progresses, exercises can incorporate balance

 © 2026 Frosh H

challenges, controlled directional changes, and simple visual-response activities. More advanced stages may include ball-related tasks, unpredictable movement instructions, and position-specific tactical situations. The progression should avoid introducing complex cognitive demands before the athlete demonstrates adequate control during simpler physical activities (Table 3).

Table 3. Progressive Training Objectives During Cognitive-Motor Rehabilitation.

Rehabilitation stage	Physical objective	Cognitive-motor activity
Early rehabilitation	Restore mobility and basic muscular function	Simple visual recognition and response tasks
Intermediate rehabilitation	Improve strength, balance, and movement stability	Controlled dual-task exercises
Advanced rehabilitation	Develop acceleration and directional control	Reactive movement and ball-related activities
Field-based rehabilitation	Restore football-specific movement capacity	Passing, scanning, and decision-making drills
Return-to-team training	Re-establish competitive movement demands	Position-specific tactical and perceptual activities

Integration of BFR and VR

Combining BFR with VR represents an emerging approach to rehabilitation. BFR may provide a physiological stimulus through low-load exercise, while VR can maintain perceptual and cognitive engagement. A proposed intervention could involve supervised low-load lower-limb exercise combined with virtual tasks requiring visual recognition, response selection, and movement coordination. The protocol must be individualized according to clinical status, limb occlusion pressure, exercise tolerance, and rehabilitation stage. BFR should be prescribed by appropriately trained professionals following relevant screening and safety procedures. The combined intervention remains a proposed application rather than an established standard of care. Its effectiveness in professional football requires direct investigation.

DISCUSSION

The available literature indicates that successful football performance involves more than the development of strength, speed, and endurance. Athletes must coordinate physical actions with continuous visual information processing, tactical interpretation, and rapid responses to changing environmental conditions. These requirements become increasingly important during high-pressure match situations, where delayed decisions or inappropriate movement responses may affect technical execution.

The relationship between cognitive function and physical performance also has important implications for rehabilitation. Following ACL reconstruction, athletes may regain acceptable muscular strength and functional movement capacity while continuing to experience difficulties involving sensorimotor regulation. Rehabilitation based exclusively on conventional physical assessments may therefore provide an incomplete picture of competitive readiness [27].

Incorporating perceptual activities into rehabilitation may help address these limitations. Progressive visual scanning, response selection, divided-attention exercises, and unpredictable movement tasks can expose athletes to cognitive demands similar to those encountered during competition. However, these activities should be introduced according to functional recovery and individual tolerance rather than applied uniformly throughout rehabilitation. BFR offers another possible strategy for maintaining muscular conditioning when heavy resistance exercise is unsuitable. Its application may support physical adaptation through low-load exercise, although differences in cuff pressure, exercise prescription, and participant characteristics influence the resulting responses. Consequently, clinical supervision and individualized programming remain essential.

VR provides an additional opportunity to develop perceptual and decision-making abilities without requiring continuous high-intensity movement. It can simulate selected match situations and allow practitioners to regulate visual complexity and task difficulty. Nevertheless, virtual performance does not necessarily reproduce the physical, emotional, and tactical pressures of actual competition. The proposed combination of VR and BFR is scientifically interesting because it brings together cognitive engagement and physiological

 © 2026 Frosh H

stimulation. However, evidence supporting their combined effectiveness in elite football remains insufficient. Controlled investigations should evaluate neuromuscular outcomes, perceptual accuracy, movement quality, training tolerance, and subsequent competitive performance [28].

Future research should also examine differences between playing positions, injury types, rehabilitation stages, and individual cognitive profiles. Longitudinal investigations involving professional footballers would help determine whether integrated interventions provide measurable benefits beyond conventional rehabilitation. Objective assessment should include physical performance, movement mechanics, decision-making accuracy, and the athlete's ability to respond to unpredictable football-specific situations.

CONCLUSION

Professional football requires the coordinated development of physical capacity, perceptual processing, and movement regulation. Rehabilitation following major injuries should extend beyond muscular recovery to address cognitive-motor adaptations that influence competitive performance. Visual scanning, selective attention, divided attention, and rapid decision-making can be incorporated progressively into football-specific conditioning. Blood flow restriction offers a possible means of maintaining muscular stimulation when high mechanical loading is unsuitable, whereas virtual reality provides opportunities to practice perceptual responses within controlled environments. Their combined application represents an emerging rehabilitation strategy, although its effectiveness in elite football has not yet been established. Individualized progression, appropriate clinical supervision, and objective functional assessment remain essential when preparing athletes for competition. Future investigations should determine the optimal integration of these methods and establish their effects on neuromuscular function, movement quality, and reinjury outcomes. Such evidence may support more comprehensive rehabilitation pathways and improve athletes' preparedness for the physical and cognitive demands of professional football.

REFERENCES

1. Bortnik L, Bruce-Low S, Burger J, Alexander J, Harper D, et al. Transitional activities in elite football: Frequency, type, effect on match outcome and the novel concept of clusters. Res Inves Sports Med. 2023;9(5):872–886.
2. Bradley PS. Setting the benchmark part 2: Contextualising the physical demands of teams in the FIFA World Cup Qatar 2022. Biol Sport. 2024;41(1):271–278.
3. Neto T, Sayer T, Theisen D, Mierau A. Functional brain plasticity associated with ACL injury: A scoping review of current evidence. Neural Plast. 2019;2019:3480512.
4. Diekfuss JA, Grooms DR, Hogg JA, Singh H, Slutsky-Ganesh AB, et al. Targeted application of motor learning theory to leverage youth neuroplasticity for enhanced injury-resistance and exercise performance: Optimal prep. J Sci Sport Exerc. 2021;3(4):17–36.
5. Chaput M, Simon J, Taberner M, Grooms DR. From control to chaos: Visual-cognitive progression during recovery from ACL reconstruction. J Orthop Sports Phys Ther. 2024;54(7):431–439.
6. Oberauer K. Working memory and attention: A conceptual analysis and review. J Cogn. 2019;2(1):36.
7. Desimone R, Duncan J. Neural mechanisms of selective visual attention. Annu Rev Neurosci. 1995;18:193–222.
8. Ward SH, Pearce A, Bennell KL, Pietrosimone B, Bryant AL. Quadriceps cortical adaptations in individuals with an anterior cruciate ligament injury. Knee. 2016;23(4):582–587.
9. Soltanabadi S, Minoonejad H, Bayattork M, Seyedahmadi M. Effect of virtual reality and augmented reality training for injury prevention and accelerating rehabilitation of anterior cruciate ligament injury in athletes: A scoping review. Asian J Sports Med. 2023;14(4):1–19.
10. Bortnik L, Bruce-Low S, Burger J, Alexander J, Harper D, et al. Physical match demands across different playing positions during transitional play and high-pressure activities in elite soccer. Biol Sport. 2024;41(2):73–82.
11. Cunniffe B, Sharma V, Cardinale M, Yellon D. Characterization of muscle oxygenation response to vascular occlusion: Implications for remote ischaemic preconditioning and physical performance. Clin Physiol Funct Imaging. 2017;37(6):785–793.
12. Arriel R, Rodrigues J, Souza H, Meireles A, Leitão LF, et al. Ischemia-reperfusion intervention: From enhancements in exercise performance to accelerated performance recovery—A systematic review and meta-analysis. Int J Environ Res Public Health. 2020;17(21):8161.
13. Bennett H, Slattery F. Effects of blood flow restriction training on aerobic capacity and performance: A systematic review. J Strength Cond Res. 2019;33(2):572–583.
14. May AK, Russell AP, Gatta PA, Warmington SA. Muscle adaptations to heavy-load and blood flow restriction resistance training methods. Front Physiol. 2022;13:837697.
15. Patterson SD, Hughes L, Warmington S, Burr J, Scott BR, et al. Blood flow restriction exercise: Considerations of methodology, application and safety. Front Physiol.

 © 2026 Frosh H

2019;10:533.

16. Kargaran A, Abedipour A, Saadatmehr Z, Yaali R, Amani-Shalamzari S, et al. Effects of dual-task training with blood flow restriction on cognitive functions, muscle quality and circulatory biomarkers in elderly women. Physiol Behav. 2021;239:113500.
17. Neal BS, McManus CJ, Bradley WJ, Leaney SF, Murray K, et al. The feasibility, safety and efficacy of lower limb garment-integrated blood flow restriction training in healthy adults. Phys Ther Sport. 2023;60:9–16.
18. Kim D, Singh H, Loenneke JP, Thiebaud RS, Fahs CA, et al. Comparative effects of vigorous-intensity and low-intensity blood flow restricted cycle training and detraining on muscle mass, strength and aerobic capacity. J Strength Cond Res. 2016;30(5):1453–1461.
19. Schiffer T, Schulte S, Sperlich B, Achtzehn S, Fricke H, et al. Lactate infusion at rest increases BDNF blood concentration in humans. Neurosci Lett. 2011;488(3):234–237.
20. Clarkson MJ, Conway L, Warmington SA. Blood flow restriction walking and physical function in older adults: A randomized control trial. J Sci Med Sport. 2017;20(12):1041–1046.
21. Clarkson MJ, Scott BR, Patterson SD, Damascento SO, Junior EP, et al. Virtual reality-based therapy associated with blood flow restriction in older adults: A proposal for integration of techniques. Front Physiol. 2022;13:958823.
22. Groot PC, Thijssen DH, Sanchez M, Ellenkamp R, Hopman MT. Ischemic preconditioning improves maximal performance in humans. Eur J Appl Physiol. 2010;108(1):141–146.
23. Novaes J, Telles LG, Monteiro ER, Araujo G, Vingren JL, et al. Ischemic preconditioning improves resistance training session performance. J Strength Cond Res. 2021;35(11):2993–2998.
24. Patterson SD, Bezodis NE, Glaister M, Pattison JR. The effect of ischemic preconditioning on repeated sprint cycling performance. Med Sci Sports Exerc. 2015;47(8):1652–1658.
25. Faure C, Limballe A, Bideau B, Kulpa R. Virtual reality to assess and train team ball sports performance: A scoping review. J Sports Sci. 2020;38(2):192–205.
26. Jia Y, Zhou X, Yang J, Fu Q. Animated VR and 360-degree VR to assess and train team sports decision-making: A scoping review. Front Psychol. 2024;15:1410132.
27. Gérin-Lajoie M, Ciombor DM, Warren WH, Aaron RK. Using ambulatory virtual environments for the assessment of functional gait impairment: A proof-of-concept study. Gait Posture. 2010;31(4):533–536.
28. Gérin-Lajoie M, Richards CL, Fung J, McFadyen BJ. Characteristics of personal space during obstacle circumvention in physical and virtual environments. Gait Posture. 2008;27(2):239–247.

 © 2026 Frosh H