

# Patterns of Self Medication Among Educated and Uneducated Population in a University Setting in Lahore Pakistan: A Cross Sectional Survey

Sehrish Khalid<sup>1,2</sup>, Ishrat Younus<sup>1,3\*</sup>, Saima Muhammad Hussain<sup>1,4</sup>, Faiza Naveed<sup>1,5</sup>  
Javaid Iqbal<sup>1</sup>

<sup>1</sup>Faculty of Pharmacy, The University of Lahore, Lahore-Pakistan.

<sup>2</sup>Sultan Bin Abdul Aziz Humanitarian City, Riyadh.

<sup>3</sup>Faculty of Pharmacy, Hamdard University, Karachi-Pakistan.

<sup>4</sup>Eli Lilly - Oncology Department, Lahore, Pakistan.

<sup>5</sup>Amin enterprises Toba Tek Singh, Pakistan.

\*ishratyounas@gmail.com

## Abstract

**Background and Objective:** Self-medication is gaining attention and is becoming a public health concern. Malpractice of self-medication is associated with risks such as misdiagnosis, overdose, prolonged duration, drug interactions and poly-pharmacy; which causes increased burden on the individual and is problematic in the elderly. This study evaluated the prevalence and patterns of self-medication among Educated and Uneducated population in a university setting in Pakistan.

**Methods:** A descriptive cross-sectional study was conducted in 100 individuals (educated: 50 and uneducated: 50). A questionnaire was filled consisting questions related to symptoms that necessitate self-medication, motive for self-medication and the sources of information about medication.

**Results:** There was no significant difference in prevalence of self-medication in educated (68%) and uneducated samples (60%). Headache (32%), fever (30%), cough (16%) and gastric problems (8%) were the common symptoms for which self-medication were sought. However, Self-mediation for headache (36% vs 28%) and fever (32% vs 28%) was significantly higher in uneducated vs educated sample. The main reasons to self-medicate among educated population vs uneducated included conditions not serious (48% vs 32%); lack of time (48% vs. 20%), whereas, 44% uneducated reported to save money as compared to 4% educated sample. The common source for self-medication in both the samples were past experience (56%) and advice from others (24%), however, advertisement as a source of self-medication prevailed in uneducated sample (20%) as compared to educated sample (8%).

**Conclusion:** Self-medication is prevalent in educated and uneducated population. The drug authorities must regulate policies to provide safe and cost effective products, in particular to safe antibiotic usage and prevent adverse effects of drugs. Standards should be applied to the provision of information and advertisement to ensure responsible self-medication.

**Keywords:** self-medication, educated, uneducated

## INTRODUCTION

Self-medication involves the use of medicinal products by the consumers to treat self-diagnosed disorders or symptoms, or the intermittent or continued use of a prescribed drug for chronic or recurrent disease or symptoms [6]. It has also been defined as obtaining and consuming medication without professional supervision, which

## **Patterns of Self Medication Among Educated and Uneducated Population in a University Setting in Lahore, Pakistan: A Cross Sectional Survey**

---

comprises of acquiring medicines without a prescription, purchasing drugs by resubmitting/ reutilizing an old prescription, taking medicines on advice of relative or others, or consuming left-over medicines already available at home[2].

Self-medication is globally practiced with varying degree. A systematic review done by Montgomery et al., self-medication prevalence rate was greater than 50%. Self-medication can result in severe health-related complications when implemented inappropriately without any regulations. For instance, resistance to antibiotics is of great concern due to frequent and inappropriate use. Resistance prolongs illnesses and hospital stays, and can even cause death, leading to costs of US\$ 4–5 billion per year in the United States of America and €9 billion per year in Europe. Moreover due to irresponsible self-medication, consumer can be at potential risks of incorrect self-diagnosis, masking underlying disease, incorrect choice of therapy, overuse/underuse, unnecessary prolonged use, risk of dependence and abuse, therapeutic duplication, polypharmacy and severe adverse effects. Consequently, it could result in an increase in drug-induced disease and in wasteful public expenditure causing increased economic burden.

Irrational use of drugs is a wide spread public health concern. The alarmingly high prevalence rates (68% in European countries), while much higher in the developing countries needs to be addressed. Research showed that the trend to practice self-medication is higher in literates as compared to illiterates. Despite this, self-medication is scarcely focused in Pakistan and insufficient measures have been taken to address this problem. One study discussed over-the-counter availability of antimicrobial agents, self-medication and patterns of resistance [5] and suggested continuous education of health care professionals to control misuse of antimicrobials. The goal of this study is to discern the attitude and practice towards self-medication between educated and uneducated population and suggest measures that can be employed to promote responsible safe self-medication.

### **METHODOLOGY**

A cross-sectional survey was conducted in University of Lahore during April 2013. The study population was adults (over 16 years old) living in Lahore City. A convenience sample of 100 participants, 50 educated and 50 uneducated was taken by approaching them. A questionnaire was developed using combination of open-ended and close-ended questions from various previously conducted similar studies. Questions included the symptoms for which medication were being used; the types of medication; the duration of use; the reasons for self-medication; the sources of information about medication. Socio-demographic characteristics involved gender, age, marital status, education, occupation. Workers in the university mainly sweeper, gardener, guards and maids were approached that constituted the uneducated population. The data collector read the questions and wrote the answers on behalf of the uneducated participant. Similarly students, lecturers and hostel mates of different disciplines comprised the educated population. Data management and analysis was done on SPSS version 18. Descriptive statistics has been applied. Variables included age, gender, type of medication, condition for which medication was used, duration, other conditions for which self-medication is practiced, source and reasons of self-medication. Frequency tables and bar charts have been analysed.

### **RESULTS**

100 participants aged between 16-70 years were approached. The mean age of educated and uneducated participants was 24 years and 40 years respectively. In our survey 48% (n= 24) comprises of female uneducated population while 52% (n = 26) were uneducated male. Among educated population, 60% (n = 30) were female and 40% male (n = 20).

There was no significant difference in prevalence of self-medication in educated (68%) and uneducated sample (60%). Few responders have found to be taking homeopathic and herbal medication as their first choice of treatment alternative to allopathic medicine. The main reason for taking homeopathic and herbal medicine reported to be a safer type of medication and easy availability of herbal products at home. The type of medication

## Patterns of Self Medication Among Educated and Uneducated Population in a University Setting in Lahore, Pakistan: A Cross Sectional Survey

was unambiguous in uneducated population because most of them were unaware of the name of medication and reported the medication was given by the retailers at medical store after being informed of their condition.

Headache (32%), fever (30%), cough (16%) and gastric problems (8%) were the common symptoms for which self-medication were sought. However, Self-mediation for headache (36% vs 28%) and fever (32% vs 28%) was significantly higher in uneducated vs educated sample. The uneducated retired guards have been observed to be taking analgesics for no reason on daily basis for mere satisfaction. Reported medicines included dispirin, panadol, bruffin, ponstan, flagyl, augmentin, arinac, cough syrups, vitamin supplements, flagyl, omeprazole and various similar brand alternatives. In other conditions pain at different body parts, lethargy and sedation were reported. One case reported gastritis but claimed NSAIDS necessary to take relief from pain. One individual mentioned eccentric behaviour for antibiotic self-medication, taking one antibiotic for one day and other antibiotic the other day without completing any of the courses. Some medications were taken daily by some of the individuals which became their habit and reported they feel lethargic if they don't take the medication.

The main reasons to self-medicate among educated population vs uneducated included conditions not serious (48% vs 32%); lack of time (48% vs 20%), whereas, 44% uneducated reported to save money as compared to 4% educated sample. One of the uneducated female reported the inaccessibility of physician. One uneducated male reported that physician usually panic the patient by creating an alarming state. One educated female reported that she usually self-medicate for a known disease as the physician is to prescribe the similar medications (Table 1).

**Table1. Reason for self-medication**

		Frequency	Percent
uneducated	lack of time	10	20.0
	save money	22	44.0
	problem not serious	16	32.0
	others	2	4.0
	Total	50	100.0
educated	lack of time	24	48.0
	save money	2	4.0
	problem not serious	24	48.0
	Total	50	100.0

The common source for self-medication in both the samples were past experience (56%) and advice from others (24%), however, advertisement as a source of self-medication prevailed in uneducated sample (20%) as compared to educated sample (8%) (Table 2).

**Table2. Source**

		Frequency	Percent
uneducated	Advertisement	10	20.0
	heard from others	12	24.0
	past experience	28	56.0
	Total	50	100.0
educated	advertisement	4	8.0
	heard from others	12	24.0
	past experience	28	56.0
	others	6	12.0
	Total	50	100.0

## **DISCUSSION**

Self-medication involves the use of medicinal products by the consumers to treat self-diagnosed disorders or symptoms, or the intermittent or continued use of a prescribed drug for chronic or recurrent disease or symptoms [6]. Easy availability of the drugs over the counter facilitates self-medication [21]. This leads to inappropriate self-medication masking the sign and symptoms of underlying disease thereby delay in diagnosis. Although there was no significant difference in prevalence of self-medication in educated and uneducated participants, the scope and duration of self-medication can be kept within safe limits by limiting the package sizes or dispensing dosages for not more than 3 days to avoid stocking of medication at home and avoid prolonged use. The community pharmacies should keep a record of their community prescription medications to avoid therapeutic duplication and polypharmacy. Government should disseminate effective education of self-medication to protect from possible harm and long-term adverse effects of medications. There should be a culture of reporting adverse drug reactions by consumers through pharmacies by giving necessary information to consumers about adverse drug reactions of products and when to report. Promotional messages through the media and the Internet should be utilized for this purpose. Advice to the consumer/patient should include a description of how to use the product without medical supervision and the circumstances in which referral for medical advice is necessary [3]. Inappropriate use of antibiotics increases antimicrobial resistance and render medicines ineffective against infectious disease. Governments should recognize and enforce the distinction between prescription and non-prescription medicines [10]. Therefore, drug authorities should restrict antibiotics as prescription only medicines.

## **CONCLUSION**

Self-medication is prevalent in educated and uneducated population. The drug authorities must regulate policies to provide safe and cost effective products, in particular to safe antibiotic usage and prevent adverse effects of drugs. Standards should be applied to the provision of information and advertisement to ensure responsible self-medication.

## **REFERENCES**

1. Hughes CM1, McElnay JC, Fleming GF; Benefits and risks of self medication. *Drug Saf.* 2001; 24(14):1027-37.
2. Hussain A, Khanum A. Self medication among university students of Islamabad, Pakistan- a preliminary study. *Southern Med Review* (2008) 1; 1:14-16
3. The Benefits and Risks of Self-medication. (WHO Drug Information Vol. 14, No. 1, 2000) Retrieved from: <http://apps.who.int/medicinedocs/en/d/Js2218e/1.1.html>
4. World Health Organization (1998): The Role of the Pharmacist in Self-Care and Self-Medication (WHO/DAP/98.13) Retrieved from: <http://apps.who.int/medicinedocs/pdf/whozip32e/whozip32e.pdf>
5. Sturm AW, van der Pol R, Smits AJ, van Hellemond FM, Mouton SW, Jamil B, et al. Over-the-counter availability of antimicrobial agents, self-medication and patterns of resistance in Karachi, Pakistan. *J Antimicrob Chemother.* 1997;39:543-7. 10.1093/jac/39.4.543
6. World Health Organization (2000): Guidelines for the Regulatory Assessment of Medicinal Products for Use in Self-Medication. Geneva: (WHO/EDM/QSM/00.1) Retrieved from: <http://apps.who.int/medicinedocs/pdf/s2218e/s2218e.pdf>
7. D Brahma, M Marak, J Wahlang. Rational Use of Drugs and Irrational Drug Combinations. *The Internet Journal of Pharmacology.* 2012 Volume 10 Number 1.

## Patterns of Self Medication Among Educated and Uneducated Population in a University Setting in Lahore, Pakistan: A Cross Sectional Survey

8. World Self - Medication Industry (WSMI). What is Self- Medication? Retrieved from: <http://www.wsmi.org/about-self-care-and-self-medication/what-is-self-medication/>
9. World Self - Medication Industry (WSMI). The story of Self-Care and Self-Mediation. 40 years of progress, 1970-2010. Retrieved from: [http://www.wsmi.org/wp-content/data/pdf/storyofselfcare\\_bdpape.pdf](http://www.wsmi.org/wp-content/data/pdf/storyofselfcare_bdpape.pdf)
10. World Medical Association (WMA) Statement on Self-Medication 2002. Retrieved from: <https://www.wma.net/policies-post/wma-statement-on-self-medication/>
11. Montgomery AJ, Bradley C, Rochfort A, Panagopoulou E. A review of self-medication in physicians and medical students. *Occup Med.* 2011;61(7):490–497.
12. Abay SM, Amelo W. Assessment of Self-Medication Practices Among Medical, Pharmacy, and Health Science Students in Gondar University, Ethiopia. *Journal of Young Pharmacists : JYP.* 2010;2(3):306-310. doi:10.4103/0975-1483.66798.
13. Shankar PR, Partha P, Shenoy N. Self-medication and non-doctor prescription practices in Pokhara valley, Western Nepal: a questionnaire-based study. *BMC Fam Pract* 2002; 3: 17.
14. Hsiao FY, Alee J, Huang W-F, Chen S-M, Cheny H-Y. Survey of medication knowledge and behaviors among college students in Taiwan. *Am J Pharm Educ* 2006; 70: 30.
15. Zafar, S., Syed, R., Waqar, S., Zubairi, A., Vaqar, T., Shaikh, M., Yousaf, W., Shahid, S., Saleem, S. (2008). Self-medication amongst university students of Karachi: prevalence, knowledge and attitudes. *Journal of the Pakistan Medical Association*, 58(4), 214-7.
16. Kumar, N., Kanchan, T., Unnikrishnan, B., Rekha, T., Mithra, P., Kulkarni, V., ... Uppal, S. (2013). Perceptions and Practices of Self-Medication among Medical Students in Coastal South India. *PLoS ONE*, 8(8), e72247. <http://doi.org/10.1371/journal.pone.0072247>
17. Joint Statement by the International Pharmaceutical Federation and The World Self-Medication Industry: Responsible Self Medication Available: [http://www.fip.org/www/uploads/database\\_file.php?id=241&table\\_id](http://www.fip.org/www/uploads/database_file.php?id=241&table_id).
18. Grigoryan, L., Haaijer-Ruskamp, F. M., Burgerhof, J. G. M., Mechtler, R., Deschepper, R., Tambic-Andrasevic, A., ... Birkin, J. (2006). Self-medication with Antimicrobial Drugs in Europe. *Emerging Infectious Diseases*, 12(3), 452–459. <http://doi.org/10.3201/eid1203.050992>
19. World Self - Medication Industry (WSMI). Guiding Principles in Self-Medication. Retrieved from: [http://www.wsmi.org/wp-content/data/pdf/GuidingPrinciplesinSelfMedication\\_I.pdf](http://www.wsmi.org/wp-content/data/pdf/GuidingPrinciplesinSelfMedication_I.pdf)
20. El Ezz NF, Ez-Elarab HS ,Knowledge, attitude and practice of medical students towards self-medication at Ain Shams University. *Egypt J Prev Med Hyg* 2011;52:196–200.
21. Yasmin, S.M.; Ashraf, J.; Tahira, M.; Shahla, Z.; Adnan, S. Self medication among university students of Karachi. *JLUMHS* 2011, 10, 102–105.

**Citation:** Sehrish Khalid, Ishrat Younus, Saima Muhammad Hussain, Faiza Naveed, Javaid Iqbal, "Patterns of Self Medication Among Educated and Uneducated Population in a University Setting in Lahore, Pakistan: A Cross Sectional Survey". *American Research Journal of Pharmacy.* 2019; 4(1): 1-5.

**Copyright © 2019 Sehrish Khalid, Ishrat Younus, Saima Muhammad Hussain, Faiza Naveed, Javaid Iqbal, This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**